

IGCSE MATHEMATICS FORMULA HANDBOOK

**Distribute this book freely. NOT FOR SALE*

Content

Natural Numbers.....	1	Congruent Triangles.....	8
Whole Numbers	1	Similar Triangles	9
Integers.....	1	Area of Similar Triangles.....	9
Rational Numbers.....	1	Polygons.....	10
Irrational Numbers	1	Similar Solids	10
Terminating Decimals	1	CIRCLE.....	10
Recurring Decimals	1	Chord of a circle	11
Significant figures	2	Tangent to a Circle	12
Decimal Places	2	Law of Indices.....	12
Standard Form	2	Solving Inequalities.....	12
Conversion Factors	2	TRIGONOMETRY.....	13
Time	2	Bearing	13
Percentages.....	3	Cartesian co-ordinates	15
Simple Interest To find the interest:	3	Distance – Time Graphs	15
Compound Interest.....	3	Speed – Time Graphs	15
Speed, Distance and Time.....	3	Velocity	15
Quadratic Equations.....	4	Acceleration	15
Expansion of algebraic expressions	4	SETS.....	16
Factorization of algebraic expressions	4	Loci and construction.....	17
Ordering	4	Vectors:.....	17
Variation	4	Column Vectors:.....	18
PHTHAGORAS' THEOREM	5	Parallel Vectors:	18
Area and Perimeter	5	Modulus of a Vector:.....	18
Surface Area and Volume	6	MATRICES:	18
(a) Angles on a straight line	7	The Inverse of a Matrix:	19
(c) Vertically opposite angles.....	7	Transformations:	19
Parallel Lines.....	7	Transformation by Matrices.....	22
Different types of triangles.....	8	STATISTICS.....	23
Angle properties of triangle	8	Probability.....	25
		Symmetry	26

NUMBER

Natural Numbers: Numbers which are used for counting purpose are called natural numbers.

Ex: 1, 2, 3, 4, 100,

Whole Numbers: Natural numbers including 0 are called Whole Numbers.

Ex: 0, 1, 2, 3, 4,

Integers: Positive natural numbers, negative natural numbers along with 0 are called integers.

Ex:, -4, -3, -2, -1, 0, 1, 2, 3, 4,

Rational Numbers: Numbers which are in the form of $\frac{p}{q}$ ($q \neq 0$) where p and q are positive or negative whole numbers are called rational numbers.

Ex: $\frac{1}{2}$, $\frac{3}{4}$, $\frac{-5}{7}$, $\frac{49}{-56}$

Irrational Numbers: Numbers like $\sqrt{2}$, π cannot be expressed as rational numbers. Such types of numbers are called as irrational numbers.

Ex: $\sqrt{5}$, $\sqrt{17}$,

Terminating Decimals

These are decimal numbers which stop after a certain number of decimal places.

For example, $7/8=0.875$, is a terminating decimal because it stops (terminates) after 3 decimal places.

Recurring Decimals

These are decimal numbers which keep repeating a digit or group of digits;

for example $137/259 = 0.528\ 957\ 528\ 957\ 528\ 957\ \dots$, is a recurring decimal. The six digits 528957 repeat in this order. Recurring decimals are written with dots over the first and last digit of the repeating digits, e.g. $0.\dot{5}28\ 95\dot{7}$

- The order of operations follows the **BODMAS** rule:

Brackets
Power **O**f
Divide
Multiply
Add
Subtract

- Even numbers:** numbers which are divisible by 2, eg, 2, 4, 6, 8, ...
- Odd numbers:** numbers which are not divisible by 2, eg, 1, 3, 5, 7, ...
- Real numbers** are made up of all possible rational and irrational numbers.
- An integer is a whole number.
- A prime number is divisible only by itself and by one (1). 1 is not a prime number. It has only two factors. 1 and the number itself.
- The exact value of rational number can be written down as the ratio of two whole numbers.
- The exact value of an irrational number cannot be written down.
- A square number is the result of multiplying a number by itself.
Ex: $1^2, 2^2, 3^2, \dots$ i.e. 1, 4, 9,
- A cube number is the result of multiplying a number by itself three times.
Ex: $1^3, 2^3, 3^3, \dots$ i.e. 1, 8, 27,
- The factors of a number are the numbers which divide exactly into two.
eg. Factors of 36
1, 2, 3, 4, 6, 9, 12, 18
- Multiples of a number are the numbers in its times table.
eg. Multiples of 6 are 6, 12, 18, 24, 30, ...

Significant figures

Example:

8064 = 8000 (correct to 1 significant figures)

8064 = 8100 (correct to 2 significant figures)

8064 = 8060 (correct to 3 significant figures)

0.00508 = 0.005 (correct to 1 significant figures)

0.00508 = 0.0051 (correct to 2 significant figures)

2.00508 = 2.01 (correct to 3 significant figures)

Decimal Places

Example

0.0647 = 0.1 (correct to 1 decimal places)

0.0647 = 0.06 (correct to 2 decimal places)

0.0647 = 0.065 (correct to 3 decimal places)

2.0647 = 2.065 (correct to 3 decimal places)

Standard Form

The number $a \times 10^n$ is in standard form when $1 \leq a < 10$ and n is a positive or negative integer.

Eg: $2400 = 2.4 \times 10^3$

Conversion Factors

Length:

1 km = 1000 m

1 m = 100 cm

1 cm = 10 mm

km means kilometer

m means meter

cm means centimeter

mm means millimeter

Volume:

1 litre = 1000 cm³

1 m³ = 1000 litres

1 kilo litre = 1000 litre

1 dozen = 12

Mass:

1 kg = 1000 gm where kg means kilogram

1 gm = 1000 mgm gm means gram

1 tonne = 1000 kg mgm means milligram

Time

1 hours = 60 minutes = 3600 seconds

1 minute = 60 seconds

1 day = 24 hours

1 year = 12 months

= 52 weeks

= 365.25 days

1 week = 7 days

1 leap year = 366 days

1 light year = 9.46×10^{12} km

Percentages

- Percent means per hundred
- To express one quantity as a percentage of another, first write the first quantity as a fraction of the second and then multiply by 100
- Profit = S.P. – C.P.
- Loss = C.P. – S.P.
- Profit percentage = $\frac{SP-CP}{CP} \times 100$
- Loss percentage = $\frac{CP-SP}{CP} \times 100$

CP = Cost Price SP = Selling Price

Simple Interest

To find the interest:

- $i = \frac{PRT}{100}$ where
 P = money invested or borrowed
 R = rate of interest per annum
 T = Period of time (in years)

To find the amount:

- $A = P + I$ where A = amount


Compound Interest

$$A = p\left(1 + \frac{r}{100}\right)^n$$

Where,
 A stands for the amount of money accruing after n year,
 P stands for the principal
 R stands for the rate per cent per annum
 n stands for the number of years for which the money is invested.

Speed, Distance and Time

- Distance = speed \times time
- Speed = $\frac{\text{distance}}{\text{time}}$
- Time = $\frac{\text{distance}}{\text{Speed}}$
- Average speed = $\frac{\text{total distance}}{\text{total time}}$
- Units of speed: km / hr, m / sec
- Units of distance: km, m
- Units of time: hr, sec
- $\text{km / hr} \times \frac{5}{18} = \text{m / sec}$
- $\text{m / sec} \times \frac{18}{5} = \text{km / hr}$



ALGEBRA

Quadratic Equations

An equation in which the highest power of the variable is 2 is called quadratic equation. Thus $ax^2 + bx + c = 0$ where a, b, c are constants and $a \neq 0$ is a general equation.

Solving quadratic equations:

We can solve quadratic equation by method of,

- a) Factorization
- b) Using the quadratic formula
- c) Completing the square

(a) Solution by factors:

Consider the equation $c \times d = 0$, where c and d are numbers. The product $c \times d$ can only be zero if either c or d (or both) is equal to zero.

i.e. c

(b) Solution by formula:

The solutions of the quadratic equation

$ax^2 + bx + c = 0$ are given by the formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

(c) Completing the square

- Make the coefficient of x^2 , i.e. $a=1$
- Bring the constant term, i.e. c to the right side of equation.
- Divide coefficient of x , i.e. by 2 and add the square i.e. $(\frac{b}{2})^2$ to both sides of the equation.
- Factorize and simplify answer

<u>Expansion of algebraic expressions</u>	<u>Factorization of algebraic expressions</u>
<ul style="list-style-type: none"> • $a(b + c) = ab + ac$ • $(a + b)^2 = a^2 + 2ab + b^2$ • $(a - b)^2 = a^2 - 2ab + b^2$ • $a^2 + b^2 = (a + b)^2 - 2ab$ • $a^2 - b^2 = (a + b)(a - b)$ 	<ul style="list-style-type: none"> • $a^2 + 2ab + b^2 = (a + b)^2$ • $a^2 - 2ab + b^2 = (a - b)^2$ • $a^2 - b^2 = (a + b)(a - b)$

Ordering

= is equal to

≠ is not equal to

> is greater than

≥ is greater than or equal to

< is less than


≤ is less than or equal to

<u>Variation</u>	
Direct Variation: y is proportional to x $y \propto x$ $y = kx$	Inverse Variation: y is inversely proportional to x $y \propto \frac{1}{x}$ $y = \frac{k}{x}$

MENSURATION

PHTHAGORAS' THEOREM

For all the **right angled** triangles “the square on the hypotenuse is equal to the sum of the squares on the other two sides”



$$c^2 = a^2 + b^2$$

$$c = \sqrt{a^2 + b^2}$$








$$b = \sqrt{c^2 - a^2}$$

$$a = \sqrt{c^2 - b^2}$$

Area and Perimeter

Figure	Diagram	Area	Perimeter
Rectangle		Area= $l \times b$	perimeter= $2(l + b)$
Square		Area = side \times side = $a \times a$	Perimeter= $4 \times \text{side}$ = $4 \times a$
Parallelogram		Area= $b \times h$ Area= $ab \sin \theta$ where a, b are sides and θ is the included angle	perimeter= $2(a + b)$
Triangle		Area= $\frac{1}{2} \times \text{base} \times \text{height}$ Area= $\frac{1}{2} ab \sin C$ = $\sqrt{s(s-a)(s-b)(s-c)}$ where $s = \frac{a+b+c}{2}$	perimeter= $a + b + c$
Trapezium		Area= $\frac{1}{2}(a + b)h$	perimeter= Sum of all sides
Circle		Area= πr^2	circumference= $2\pi r$
Semicircle		Area= $\frac{1}{2}\pi r^2$	perimeter= $\frac{1}{2}\pi d + d$
Sector		Area= $\pi r^2 \times \frac{\theta}{360}$	length of an arc= $2\pi r \times \frac{\theta}{360}$


Surface Area and Volume

Figure	Diagram	Surface Area	Volume
Cylinder		curved surface area= $2\pi r h$ total surface area= $2\pi r(h + r)$	Volume= $\pi r^2 h$
Cone		curved surface area= $\pi r l$ where $l = \sqrt{r^2 + h^2}$ total surface area= $\pi r(l + r)$	Volume= $\frac{1}{3} \pi r^2 h$
Sphere		Surface area= $4\pi r^2$	Volume= $\frac{4}{3} \pi r^3$
Pyramid		Base area+area of the shapes in the sides	Volume= $\frac{1}{3}$ x base area x perpendicular height
Cuboid		Surface area= $2(lb + bh + lh)$	Volume= $l \times b \times h$
Cube		Surface area= $6l^2$	Volume= l^3
Hemisphere		Curved surface area= $2\pi r^2$	Volume= $\frac{2}{3} \pi r^3$


GEOMETRY

(a) Angles on a straight line

The angles on a straight line add up to 180° .



(b) Angle at a point



The angles at a point add up to 360° .


$$a + b + c + d = 360^\circ$$

(c) Vertically opposite angles

If two straight lines intersect, then


$$a = c$$

$$b = d \text{ (Vert. opp. } \angle\text{s)}$$



Parallel Lines

When lines never meet, no matter how far they are extended, they are said to be parallel.




- Vertically opposite angles are equal.
 $a = c$; $b = d$; $p = s$ and $q = r$
- **Corresponding angles** are equal.
 $a = q$; $b = p$; $c = r$ and $d = s$
- **Alternate angles** are equal.
 $c = q$ and $d = p$
- Sum of the angles of a triangle is 180°
- Sum of the angles of a quadrilateral is 360°


Types of angles

Given an angle, if


$\theta < 90^\circ$, then θ is an acute angle



$90^\circ < \theta < 180^\circ$, then θ is an obtuse angle



$180^\circ < \theta < 360^\circ$, then θ is a reflex angle




Triangles

Different types of triangles

1. An isosceles triangle has 2 sides and 2 angles the same.


$$AB=AC$$

$$ABC=BCA$$




2. An equilateral triangle has 3 sides and 3 angles the same.

$$AB = BC = CA \text{ and } ABC = BCA = CAB$$



3. A triangle in which one angle is a right angle is called the right angle triangle.


$$ABC = 90^\circ$$



Angle properties of triangle



- The sum of the angles of a triangle is equal to 180° .
- In every triangle, the greatest angle is opposite to the longest side. The smallest angle is opposite to the shortest side.
- Exterior angle is equal to the sum of the opposite interior angles.

$$x = a + b$$



Congruent Triangles

Two triangles are said to be congruent if they are equal in every aspect.



$$AB = XY$$

$$BC = YZ$$

$$AC = XZ$$


$$\angle a = \angle x$$

$$\angle b = \angle y$$

$$\angle c = \angle z$$


Similar Triangles

If two triangles are similar then they have a pair of corresponding equal angles and the three ratios of corresponding sides are equal.




$\angle a = \angle x; \angle b = \angle y$ and $\angle c = \angle z$ $\frac{AB}{XY} = \frac{BC}{YZ} = \frac{AC}{XZ}$

If you can show that one of the following conditions is true for two triangles, then the two triangles are similar.



- i) The angles of one triangle are equal to the corresponding angles of the other triangle.




$\triangle ABC$ is similar to $\triangle XYZ$ because $\angle a = \angle x; \angle b = \angle y$ and $\angle c = \angle z$

- ii) The ratio of corresponding sides is equal.

If $\frac{PQ}{DE} = \frac{PR}{DF} = \frac{QR}{EF}$ then $\triangle PQR$ is similar to $\triangle DEF$


- iii) The ratios of the corresponding side are equal and the angles between them are equal.



$\triangle PQR$ is similar to $\triangle XYZ$ (if, for eg: $\angle P = \angle X$ and $\frac{PQ}{XY} = \frac{PR}{XZ}$)

Area of Similar Triangles

The ratio of the areas of similar triangles is equal to the ratio of the square on corresponding sides.



$\frac{\text{area of } \triangle ABC}{\text{area of } \triangle PQR} = \frac{AB^2}{PQ^2} = \frac{BC^2}{QR^2} = \frac{AC^2}{PR^2}$

Polygons

- The exterior angles of a polygon add up to 360° .
- The sum of the interior angle of a polygon is $(n - 2) \times 180^\circ$ where n is the number of sides of the polygon.
- A regular polygon has equal sides and equal angles.
- If the polygon is regular and has n sides, then each exterior angle = $\frac{360}{n}$
-

3 sides = triangle	4 sides = quadrilateral	5 sides = pentagon
6 sides = hexagon	7 sides = heptagon	8 sides = octagon
9 sides = nonagon	10 sides = decagon	

Similar Solids

If two objects are similar and the ratio of corresponding sides is k , then

the ratio of their areas is k^2 .

the ratio of their volumes is k^3 .

Length


$$\frac{l_1}{l_2} = \frac{r_1}{r_2} = \frac{h_1}{h_2}$$

Area

$$\frac{A_1}{A_2} = \frac{r_1}{r_2} = \frac{h_1}{h_2}$$


Volume

$$\frac{l_1}{l_2} = \frac{r_1}{r_2} = \frac{h_1}{h_2}$$




CIRCLE


- The angle subtended by an arc at the centre is twice the angle subtended at the circumference




- Angles subtended by an arc in the same segment of a circle are equal.




- The angle in a semi-circle is a right angle. [or if a triangle is inscribed in a semi-circle the angle opposite the diameter is a right angle]. $\angle APB = 90^\circ$



- Opposite angles of a cyclic quadrilateral add up to 180° (supplementary). The corners touch the circle. $A+C=180^\circ$, $B+D=180^\circ$



- The exterior angle of a cyclic quadrilateral is equal to the interior opposite angle. ($b = p$)




Chord of a circle

A line joining two points on a circle is called a **chord**.

The area of a circle cut off by a chord is called a **segment**.

AXB is **the minor arc** and AYB is **the major arc**.

- The line from the centre of a circle to the mid-point of a chord bisects the chord at right angles.
- The line from the centre of a circle to the mid-point of a chord bisects the angle subtended by the chord at the centre of the circle.



Tangent to a Circle

- The angle between a tangent and the radius drawn to the point of contact is 90° .

$$\angle ABO = 90^\circ$$

- From any point outside a circle just two tangents to the circle may be drawn and they are of equal length.





$$TA = TB$$

- Alternate Segment Theorem**

The angle between a tangent and a chord through the point of contact is equal to the angle subtended by the chord in the alternate segment.

$$\angle QAB = \angle ACB \quad (p = q)$$

- $TC^2 = AC \times BC$



Law of Indices

- $a^m \times a^n = a^{m+n}$
- $a^m \div a^n = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$
- $a^{-n} = \frac{1}{a^n}$
- $(a \times b)^m = a^m \times b^m$
- $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$
- $(\sqrt[n]{a})^m = a^{m/n}$
- $\sqrt{a} \times \sqrt{b} = \sqrt{a \times b}$
- $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$
- $(\sqrt{a})^2 = a$

Solving Inequalities

When we multiply or divide by a negative number the inequality is reversed.

Eg: $4 > -2$

By multiplying by -2 [$4(-2) < (-2)(-2)$]

$-8 < +4$

TRIGONOMETRY


Let ABC be a right angled triangle, where $\angle B = 90^\circ$

$$\sin \theta = \frac{\text{Opposite Side}}{\text{Hypotenuse}} = \frac{O}{H}$$

$$\cos \theta = \frac{\text{Adjacent Side}}{\text{Hypotenuse}} = \frac{A}{H}$$


$$\tan \theta = \frac{\text{Opposite Side}}{\text{Adjacent Side}} = \frac{O}{A}$$

SOH CAH TOA



Sine Rule:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$



Cosine Rule:

To find the length of a side:

- $a^2 = b^2 + c^2 - 2bc \cos A$
- $b^2 = a^2 + c^2 - 2ac \cos B$
- $c^2 = a^2 + b^2 - 2ab \cos C$

To find area:

$$\frac{1}{2} ab \sin C$$

To find an angle when all the three sides are given:


- $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$
- $\cos B = \frac{a^2 + c^2 - b^2}{2ac}$
- $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$

Bearing


The bearing of a point B from another point A is;

- a) an angle measured from the north at A.
- b) In a clockwise direction.
- c) Written as three-figure number (i.e. from 000° to 360°)

Eg: The bearing of B from A is 050° .



Distance and Gradient



Distance Between Point A and C=

$$\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

Gradient of line AC.


$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Or


Gradient of a line,

$$m = \left(\frac{y - \text{intercept}}{x - \text{intercept}} \right)$$

Parallel Lines



Perpendicular Lines



When 2 lines are parallel,


$$m_1 = m_2$$

When 2 lines are perpendicular to each other.


$$m_1 \times m_2 = -1$$

m_1 = gradient of line 1
 m_2 = gradient of line 2

Midpoint



A point dividing a segment of a line



Midpoint,

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

A point dividing a segment of a line

$$P = \left(\frac{nx_1 + mx_2}{m + n}, \frac{ny_1 + my_2}{m + n} \right)$$



Cartesian co-ordinates

Gradient and equation of a straight line

The gradient of the straight line joining any two given points $A(x_1, y_1)$ and $B(x_2, y_2)$ is;

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

The gradient/ intercept form of the equation of a straight line is $y = mx + c$, where m =gradient and c =intercept on y-axis.




The midpoint of the line joining two points $A(x_1, y_1)$ and $B(x_2, y_2)$ is; $M = \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$

The distance between two points $A(x_1, y_1)$ and $B(x_2, y_2)$ is; $AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Parallel lines have the same gradient.

In a graph, gradient = $\frac{\text{Vertical height}}{\text{Horizontal height}}$ or $\frac{y}{x}$



Distance – Time Graphs

From O to A: Uniform speed

From B to C: uniform speed

From A to B: Stationery (speed = 0)

The **gradient** of the graph of a distance-time graph gives the **speed** of the moving body.

Speed – Time Graphs


From O to A: Uniform speed

From A to B: Constant speed (acceleration = 0)

From B to C: Uniform deceleration/ retardation

The **area** under a speed-time graph represents the **distance travelled**.

The **gradient** of the graph is the **acceleration**. If the acceleration is negative, it is called deceleration or retardation. (The moving body is slowing down.)



Velocity

Velocity is the rate of change of distance with respect to the time.

Acceleration

Acceleration is the rate of change of velocity with respect to time.


SETS

Notations

- ξ = universal set
- \cup (*union*) = all the elements
- \cap (*intersection*) = common elements
- \emptyset = empty set
- \in = belongs to
- \notin = does not belongs to
- \subseteq = Subset
- A' = compliment of A (i.e. the elements of ξ – the elements of A)
- $n(A)$ = the number of elements in A.
- De Morgan's Laws: $(A \cup B)' = (A' \cap B')$
 $(A \cap B)' = (A' \cup B')$


Subset \subseteq

$B \subseteq A$ means every elements of set B is also an element of set A.




Proper subset \subset

$B \subset A$ means every element of B is an element of set A but $B \neq A$.




Disjoint sets

Disjoint set do not have any element is common. If A and B are disjoint sets, then $A \cap B = \emptyset$




Intersection \cap

$A \cap B$ is the set of elements which are in A and also in B



Union \cup

$A \cup B$ is the set of elements in either A, B or both A and B.



Complement

The complement of A, written as A' refers to the elements in ξ but not in A.




Loci and construction


The locus of a point is a set of points satisfying a given set of conditions.

(a) **Locus of points at a distance x from a given point, O .**

Locus: *The circumference of a circle centre O , radius x .*




(b) **Locus of a points at a distance x from a straight line AB**



Locus: *A pair of parallel lines to the given line AB .*


(c) **Locus of points equidistance between 2 points.**

Locus: *Perpendicular bisector of the two points.*




(d) **Locus of points equidistant from two given lines AB and AC**

Locus: *Angle bisector of $\angle BAC$*



Vectors:


- A vector quantity has both magnitude and direction.
- Vectors a and b represented by the line segments can be added using the parallelogram rule or the nose- to- tail method.
- A scalar quantity has a magnitude but no direction. Ordinary numbers are scalars.
- The negative sign reverses the direction of the vector.
- The result of $\mathbf{a} - \mathbf{b}$ is $\mathbf{a} + \mathbf{-b}$
i.e. subtracting \mathbf{b} is equivalent to adding the negative of \mathbf{b} .



Addition and subtraction of vectors

$$\vec{OA} + \vec{AC} = \vec{OC} \text{ (Triangular law of addition)}$$

$$\vec{OB} + \vec{OA} = \vec{OC} \text{ (parallelogram law of addition)}$$



Column Vectors:

The top number is the horizontal component and the bottom number is the vertical component

$$\begin{pmatrix} x \\ y \end{pmatrix}$$

Parallel Vectors:

Vectors are parallel if they have the same direction. Both components of one vector must be in the same ratio to the corresponding components of the parallel vector.

In general the vector $k\begin{pmatrix} a \\ b \end{pmatrix}$ is parallel to $\begin{pmatrix} a \\ b \end{pmatrix}$

Modulus of a Vector:

The modulus of a vector a , is written as $|a|$ and represents the length (or magnitude) of the vector.

In general, if $x = \begin{pmatrix} m \\ n \end{pmatrix}$, $|x| = \sqrt{m^2 + n^2}$

MATRICES:

Addition and Subtraction:

Matrices of the same order are added (or subtracted) by adding (or subtracting) the corresponding elements in each matrix.

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} p & q \\ r & s \end{pmatrix} = \begin{pmatrix} a+p & b+q \\ c+r & d+s \end{pmatrix}$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} - \begin{pmatrix} p & q \\ r & s \end{pmatrix} = \begin{pmatrix} a-p & b-q \\ c-r & d-s \end{pmatrix}$$

Multiplication by a Number:

Each element of a matrix is multiplied by the multiplying number.

$$k \times \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} ka & kb \\ kc & kd \end{pmatrix}$$

Multiplication by another Matrix:

Matrices may be multiplied only if they are compatible. The number of columns in the left-hand matrix must equal the number of rows in the right-hand matrix.

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \times \begin{pmatrix} p & q \\ r & s \end{pmatrix} = \begin{pmatrix} ap + br & aq + bs \\ cp + dr & cq + ds \end{pmatrix}$$

In matrices A^2 means $A \times A$. [you must multiply the matrices together]

The Inverse of a Matrix:

$$\text{if } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ then } A^{-1} = \frac{1}{(ad-bc)} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

$AA^{-1} = A^{-1}A = I$ where I is the identity matrix.

The number $(ad - bc)$ is called the **determinant** of the matrix and is written as $|A|$

If $|A| = 0$, then the **matrix has no inverse**.

Multiplying by the inverse of a matrix gives the same result as dividing by the matrix.

e.g. if $AB = C$

$$A^{-1}AB = A^{-1}C$$

$$B = A^{-1}C$$

$$\text{if } C = \begin{pmatrix} x \\ y \end{pmatrix} \text{ and } D = \begin{pmatrix} r \\ s \end{pmatrix} \text{ then } C + D = \begin{pmatrix} x+r \\ y+s \end{pmatrix}$$

Transformations:

a) Reflection:

When describing a reflection, the position of the mirror line is essential.

b) Rotation:

To describe a rotation, the centre of rotation, the angle of rotation and the direction of rotation are required.

A clockwise rotation is negative and an anticlockwise rotation is positive.

>> (angle) (Direction)rotation about (centre)

c) Translation:

When describing a translation it is necessary to give the translation vector $\begin{pmatrix} x \\ y \end{pmatrix}$

- +x represents movement to the right
- -x represents movement to the left
- +y represents movement to the top
- -y represents movement to the bottom.

>> Translation by the column vector

d) Enlargement:

To describe an enlargement, state;

- The scale factor, K
- The centre of enlargement (the invariant point)

$$\text{Scale factor} = \frac{\text{length of the image}}{\text{length of the object}}$$

>> Enlargement by the scale factor ---centre---

If $K > 0$, both the object and the image lie on the same side of the centre of enlargement.

If $K < 0$, the object and the image lie on opposite side of the centre of enlargement.

If the scale factor lies between 0 and 1, then the resulting image is smaller than the object.

[although the image is smaller than the object, the transformation is still known as an enlargement]

$$\text{Area of image} = K^2 \text{ area of object}$$

Repeated Transformations:

$XT(P)$ means 'perform transformation T on P and then perform X on the image.'

$XX(P)$ may be written $X^2(P)$.

Inverse Transformations:

The inverse of a transformation is the transformation which takes the image back to the object.

If translation T has a vector $\begin{pmatrix} x \\ y \end{pmatrix}$, then the translation which has the opposite effect has vector $\begin{pmatrix} -x \\ -y \end{pmatrix}$

This is written as T^{-1} .

If rotation R denotes 90° clockwise rotation about $(0, 0)$, then R^{-1} denotes 90° anticlockwise rotation about $(0, 0)$.

For all reflections, the inverse is the same reflection.


Base vectors

The base vectors are considered as $I = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $J = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

The columns of a matrix give us the images of I and J after the transformation.

Shear:

$$\text{Shear factor} = \frac{\text{Distance a point moves due to the shear}}{\text{Perpendicular distance of the point from the fixed line}} = \frac{a}{b}$$




[The shear factor will be the same calculated from any point on the object with the exception of those on the invariant line]

$$\text{Area of image} = \text{Area of object}$$

Stretch:

To describe a stretch, state;

- i. the stretch factor, p
- ii. the invariant line,
- iii. the direction of the stretch
(always perpendicular to the invariant line)



Perpendicular distance of C from AB

$$\text{Scale factor} = \frac{\text{Perpendicular distance of } C' \text{ from } AB}{\text{Perpendicular distance of } C \text{ from } AB}$$

Where, P is the stretch factor

$$\text{Area of image} = p \times \text{Area of object}$$

Transformation by Matrices

Reflection

Matrix	Transformation
$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Reflection in the x-axis
$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	Reflection in the y-axis
$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	Reflection in the line $y=x$
$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$	Reflection in the line $y=-x$

Rotation

Matrix	Angle	Direction	Centre
$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	90°	Anticlockwise	(0, 0)
$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	90°	Clockwise	(0, 0)
$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	180°	Clockwise/anticlockwise	(0, 0)

Enlargement

$\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$ where k = scale factor and centre of enlargement = (0, 0)

Stretch

Matrix	Stretch factor	Invariant line	Direction
$\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}$	k	y - axis	Parallel to x - axis
$\begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}$	k	x - axis	Parallel to y - axis

Shear

Matrix	Stretch factor	Invariant line	Direction
$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$	k	x - axis	Parallel to x - axis
$\begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix}$	k	y - axis	Parallel to y - axis


Matrix Transformation

$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, Identity Matrix, no transformation.


$B = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$, Reflection about y-axis.

$C = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, Reflection about x-axis.


$D = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, Reflection about $y=x$.



$E = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$, Reflection about $y=-x$.



$F = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, Rotation for 90° anticlockwise  about origin $(0, 0)$.

$G = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, Rotation for 90° clockwise  about origin $(0, 0)$.

$H = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$, Rotation for 180° about origin $(0, 0)$.

STATISTICS

Bar Graph:

A bar chart makes numerical information easy to see by showing it in a pictorial form. The width of the bar has no significance. The length of each bar represents the quantity.

Pie Diagram:

The information is displayed using sectors of a circle.

Histograms:

A histogram displays the frequency of either continuous or grouped discrete data in the form of bars.

The bars are joined together.

The bars can be of varying width.

The frequency of the data is represented by the area of the bar and not the height.

[When class intervals are different it is the area of the bar which represents the frequency not the height]. Instead of frequency being plotted on the vertical axis, frequency density is plotted.

$$\text{Frequency density} = \frac{\text{frequency}}{\text{class width}}$$

Mean:

The mean of a series of numbers is obtained by adding the numbers and dividing the result by the number of numbers.

Mean = $\frac{\Sigma fx}{\Sigma f}$ where Σfx means 'the sum of the products'

i.e. Σ (the number \times frequency)
and Σf means 'the sum of the frequencies'.

Median:

The median of a series of numbers is obtained by arranging the numbers in ascending order and then choosing the number in the 'middle'. If there are two 'middle' numbers the median is the average (mean) of these two numbers.

Mode:

The mode of a series of numbers is simply the number which occurs most often.


Frequency tables:

A frequency table shows a number x such as a score or a mark, against the frequency f or number of times that x occurs.

Cumulative frequency:

Cumulative frequency is the total frequency up to a given point.

Cumulative frequency Curve:



A cumulative frequency curve shows the median at the 50th percentile of the cumulative frequency. The value at the 25th percentile is known as the lower quartile and that at the 75th percentile as the upper quartile.

A measure of the spread or dispersion of the data is given by the inter-quartile range where **Inter-quartile range = upper quartile – lower quartile.**

Probability

- Probability is the study of chance, or the likelihood of an event happening.
- Probability of an event = $\frac{\text{number of favourable outcomes}}{\text{Total number of equally likely outcome}}$
- If the probability = 0 it implies the event is impossible
- If the probability = 1 it implies the event is certain to happen.
- All probabilities lie between 0 and 1.
- Probabilities are written using fractions or decimals.

Exclusive events:

wo events are exclusive if they cannot occur at the same time.

The OR Rule:

For exclusive events A and B

$$p(A \text{ or } B) = p(A) + p(B)$$

Independent events:

Two events are independent if the occurrence of one even is unaffected by the occurrence of the other.

The AND Rule:

$$p(A \text{ and } B) = p(A) \times p(B)$$

where $p(A)$ =probability of A occurring

where $p(B)$ =probability of B occurring

Tree diagrams:


A tree diagram is a diagram used to represent probabilities when two or more events are combined.

Symmetry

- A line of symmetry divides a two-dimensional shape into two congruent (identical) shapes.
- A plane of symmetry divides a three-dimensional shape into two congruent solid shapes.
- A two-dimensional shape has rotational symmetry if, when rotated about a central point, it fits its outline. The number of times it fits its outline during a complete revolution is called the order of rotational symmetry.

Shape	Number of Lines of Symmetry	Order of Rotational Symmetry
Square	4	4
Rectangle	2	2
Parallelogram	0	2
Rhombus	2	2
Trapezium	0	1
Kite	1	1
Equilateral Triangle	3	3
Regular Hexagon	6	6


Terminology:



Length and Area

				<p>$r = \text{radius}$ $A = \text{area}$ $s = \text{arc length}$ $\theta = \text{angle in radian (a math)}$ $l = \text{length of chord}$ To convert degree to radian: $\theta \times \frac{\pi}{180^\circ}$</p>
Arc Length: $s = r\theta$	Length of chord: $l = 2r \sin \frac{\theta}{2}$	Area of Sector: $A = \frac{1}{2}r^2\theta$	Area of Triangle: $A = \frac{1}{2}r^2 \sin\theta$	Area of Segment: $A = \frac{1}{2}r^2(\theta - \sin\theta)$


Arc Length



$$\text{Arc length} = \frac{\theta}{360} \times 2\pi r$$

*Angle in degree

Area of a sector



$$\text{Sector of area} = \frac{\theta}{360} \times \pi r^2$$

*Angle in degree